



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bc

Search PubMed

for

Go Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

☐ 1: J Clin Pharmacol. 2002 May;42(5):547-57.

Related Articles, Link

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

Full text article at
www.jclinpharm.org

Clinical pharmacokinetics of verteporfin.

Houle JM, Strong A.

QLT Inc., Vancouver, BC, Canada.

Verteporfin, a benzoporphyrin derivative, is the first photo-sensitive (light-activated) drug to be proven effective in treating certain types of choroidal neovascularization (CNV) secondary to age-related macular degeneration (AMD). The pharmacokinetics of light-activated drugs are central to their safety and efficacy. Forty healthy Caucasian volunteers, 24 healthy Japanese volunteers, 9 patients with mild hepatic dysfunction, 69 patients with CNV due to AMD, and 21 patients with skin cancer were infused with verteporfin 3 to 20 mg/m² of body surface area over 1.5 to 45 minutes. Verteporfin regioisomers and the metabolite benzoporphyrin derivative diacid (BPD-DA) were quantified by validated methods of liquid chromatography and capillary electrophoresis with laser-induced fluorescence. C_{max} of verteporfin occurred at the end of the infusion and was proportional to the dose and rate of infusion. The extent of formation of the metabolite BPD-DA was less than 10%, based on the AUC ratio. Renal elimination was minimal (< 0.01% of the dose). All groups studied had similar pharmacokinetics, which were biexponential with distribution in the first 1 to 3 hours and elimination t_(1/2) of 5 to 6 hours. No significant differences were observed between Japanese and Caucasian volunteers or between men and women. Patients older than 65 years had a slightly higher average C_{max} than patients younger than 65 years (1.14 vs. 1.03 microg/ml, p = 0.066), but the ranges of the two age groups overlapped. Verteporfin has a short half-life and is rapidly eliminated in the bile, mainly as unchanged drug. Based on pharmacokinetic data, dose adjustments are not required for age, gender, race, or mild hepatic or renal impairment. The rapid elimination of verteporfin shows that the period of skin photosensitivity is unlikely to persist after 24 to 48 hours.

Publication Types:

- Clinical Trial
- Multicenter Study
- Randomized Controlled Trial

PMID: 12017349 [PubMed - indexed for MEDLINE]

Display	Abstract	Show: 20	Sort	Send to	Text
---------	----------	----------	------	---------	------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

Sep 4 2003 10:00-